


PREPUBLICACIONES DEL DEPARTAMENTO DE ÁLGEBRA  
DE LA UNIVERSIDAD DE SEVILLA

On the dimension of discrete valuations of  $k((X_1, \dots, X_n))$

Miguel A. Olalla-Acosta

Prepublicación nº 2 (Febrero-2000)



Departamento de Álgebra. Universidad de Sevilla



# ON THE DIMENSION OF DISCRETE VALUATIONS OF $k((X_1, \dots, X_n))$

MIGUEL ÁNGEL OLALLA ACOSTA

ABSTRACT. Let  $v$  be a rank-one discrete valuation of the field  $k((X_1, \dots, X_n))$ . We know, after [1], that if  $n = 2$  then the dimension of  $v$  is 1 and if  $v$  is the usual order function over  $k((X_1, \dots, X_n))$  its dimension is  $n - 1$ . In this paper we prove that, in the general case, the dimension of a rank-one discrete valuation can be any number between 1 and  $n - 1$ .

## TERMINOLOGY AND PRELIMINARIES

Let  $k$  be an algebraically closed field of characteristic 0,  $R_n$  the ring  $k[[X_1, \dots, X_n]]$ ,  $M_n = (X_1, \dots, X_n)$  the maximal ideal and  $K_n = k((X_1, \dots, X_n))$  the quotient field. Let  $v$  be a rank-one discrete valuation of  $K_n|k$ ,  $R_v$  its valuation ring,  $\mathfrak{m}_v$  its maximal ideal and  $\Delta_v$  its residual field of  $v$ . The center of  $v$  in  $R_n$  is  $\mathfrak{m}_v \cap R_n$ . Throughout this paper “discrete valuation of  $K_n|k$ ” will mean “rank-one discrete valuation of  $K_n|k$  whose center in  $R_n$  be the maximal ideal  $M_n$ ”. The dimension of  $v$  is the transcendence degree of  $\Delta_v$  over  $k$ . We shall suppose that the group of  $v$  is  $\mathbb{Z}$ .

Let  $\widehat{K}_n$  be the completion of  $K_n$  with respect to  $v$ ,  $\widehat{v}$  the extension of  $v$  to  $\widehat{K}_n$ ,  $R_{\widehat{v}}$ ,  $\mathfrak{m}_{\widehat{v}}$  and  $\Delta_{\widehat{v}}$  the ring, maximal ideal and the residual field of  $\widehat{v}$ , respectively. We know that  $\Delta_v$  and  $\Delta_{\widehat{v}}$  are isomorphic. Let  $\sigma : \Delta_{\widehat{v}} \rightarrow R_{\widehat{v}}$  be a  $k$ -section of the natural homomorphism  $R_{\widehat{v}} \rightarrow \Delta_{\widehat{v}}$ ,  $\theta \in R_{\widehat{v}}$  an element of value 1 and  $t$  a variable. We consider the  $k$ -isomorphism

$$\Phi = \Phi_{\sigma, \theta} : \Delta_{\widehat{v}}[[t]] \rightarrow R_{\widehat{v}}$$

given by

$$\Phi \left( \sum \alpha_i t^i \right) = \sum \sigma(\alpha_i) \theta^i,$$

and denote also by  $\Phi$  its extension to the quotient fields. We have then a  $k$ -isomorphism  $\Phi^{-1}$  which, composed with the order function on  $\Delta_{\widehat{v}}((t))$ , gives the valuation  $\widehat{v}$ . This is the situation we shall consider throughout all this paper, and we will freely use these notations and well-known results ([3]) without new explicit references.

1. THE DIMENSION OF  $v$ 

Fix a number  $m$  between 1 and  $n-1$ . We are featuring a constructive method in order to obtain examples of valuations with dimension  $m$ .

Let us consider the following homomorphism:

$$\begin{aligned} \varphi : k[[X_1, \dots, X_n]] &\longrightarrow \overline{k(u)}[[t]] \\ X_1 &\longmapsto t \\ X_2 &\longmapsto ut \\ X_i &\longmapsto \sum_{j \geq 1} u^{1/p_i^j} t^j \end{aligned}$$

where  $\overline{k(u)}$  stands for the algebraic closure of  $k(u)$  and  $2 < p_3 < \dots < p_n$  are prime numbers.

**Lemma 1.** *The homomorphism  $\varphi$  is one to one.*

*Proof.* Let us take the fields  $K_2 = k(u)$  and

$$K_i = k(u, \{u^{1/p_3^j}, j \geq 1\}, \dots, \{u^{1/p_i^j}, j \geq 1\})$$

for all  $i \geq 3$ .

Let us suppose that  $\varphi$  is not one to one, then  $\ker(\varphi) \neq \{0\}$ . So let  $f$  be a non-zero element of  $M = (X_1, \dots, X_n)$  such that  $f \in \ker(\varphi)$ . Let  $m$  be the higher index such that  $f \in k[[X_1, \dots, X_m]]$ .

If  $m = 1$  or  $2$ , trivially we have a contradiction.

If  $m = 3$ , let us take

$$\bar{f} = f(\varphi(X_1), \varphi(X_2), X_3) \in K_2[[t, X_3]],$$

and consider the homomorphism

$$\begin{aligned} \psi : K_2[[t, X_3]] &\longrightarrow \overline{k(u)}[[t]] \\ t &\longmapsto t \\ X_3 &\longmapsto \sum_{j \geq 1} u^{1/p_3^j} t^j. \end{aligned}$$

We know that  $\bar{f} \in \ker(\psi)$  and this kernel is a prime ideal because  $\psi$  is an homomorphism between integral domains. We can write  $\bar{f} = t^r g$ , with  $r \geq 0$  and  $t$  doesn't divide to  $g$ . This forces  $g$  to have some non-trivial terms in  $X_3$ . Let  $s > 0$  be the minimum such that  $\alpha X_3^s$  is one of these terms. By the Weierstrass preparation theorem we have  $g = U g'$ , where  $U(t, X_3)$  is a unit and

$$g' = X_3^s + a_1(t) X_3^{s-1} + \dots + a_s(t).$$

Since  $U$  is a unit,  $g' \in \ker(\psi)$  and

$$\psi(g') = g' \left( t, \sum_{j \geq 1} u^{1/p_3^j} t^j \right) = 0.$$

This leads to a contradiction because the roots of  $g'$  are in  $K_2[[t^{1/q}]]$ , with  $q \in \mathbb{Z}$ , by the Puiseux theorem.

If  $m > 3$  let us take

$$\bar{f} = f(\varphi(X_1), \dots, \varphi(X_{m-1}), X_m) \in K_{m-1}[[t, X_m]]$$

and consider the homomorphism

$$\begin{aligned} \psi : K_{m-1}[[t, X_m]] &\longrightarrow \overline{k(u)}[[t]] \\ t &\longmapsto t \\ X_m &\longmapsto \sum_{j \geq 1} u^{1/p_m^j} t^j. \end{aligned}$$

As in the previous case we can write  $\bar{f} = t^r h$ , where  $h \in \ker(\psi)$ . So we have  $h = U h'$ , where  $U(t, X_m)$  is a unit and

$$h' = X_m^r + b_1(t) X_m^{r-1} + \dots + b_r(t) \in \ker(\psi),$$

so

$$h' \left( t, \sum_{j \geq 1} u^{1/p_m^j} t^j \right) = 0.$$

But this is again a contradiction by the Puiseux theorem: since  $\ker(\psi)$  is a prime ideal, we can suppose that  $h'$  is an irreducible element of the ring  $K_{m-1}[[t]][X_m]$ . In this situation the Puiseux theorem says that to obtain the coefficients of a root of  $h' = 0$ , like a Puiseux series in  $t$  with coefficients in  $k(u)$ , we have to resolve a *finite number* of algebraic equations of degree greater than 1 in  $K_{m-1}$ . Inside  $K_{m-1}$  we can not obtain  $u^{1/p_m}$  and, with a finite number of algebraic equations, we can obtain a *finite number* of powers of  $u^{1/p_m^j}$  but not all. So this proves the lemma.  $\square$

We shall extend to the quotient fields this injective homomorphism for giving an example of a rank-one discrete valuation of  $k((X_1, \dots, X_n))$  of dimension 1.

**Lemma 2.** *There exists a rank-one discrete valuation of  $k((X_1, \dots, X_n))$  of dimension 1.*

*Proof.* We know that the homomorphism

$$\varphi : k((X_1, \dots, X_n)) \rightarrow \overline{k(u)}((t))$$

previously defined is one to one, so we can take the valuation  $v = \nu \circ \varphi$ , where  $\nu$  is the usual order function over  $\overline{k(u)}((t))$  in  $t$ . Let  $\alpha$  be the residue  $X_2/X_1 + \mathfrak{m}_v \in \Delta_v$ . Hence, to obtain the lemma we have to prove that  $\alpha \notin k$  and  $\Delta_v$  is an algebraic extension of  $k(\alpha)$ .

Let us suppose that  $\alpha \in k$ . Then there must exist  $a \in k$  such that  $X_2/X_1 + \mathfrak{m}_v = a + \mathfrak{m}_v$ , so

$$\frac{X_2 - aX_1}{X_1} \in \mathfrak{m}_v.$$

This means that  $v(X_2 - aX_1) > 1$ . On the other side we have

$$\varphi(X_2 - aX_1) = (u - a)t,$$

so  $v(X_2 - aX_1) = 1$  and we have a contradiction. Hence  $\alpha \notin k$ .

Let us prove that  $\Delta_v$  is an algebraic extension of  $k(\alpha)$ . We can consider each element of  $k[[X_1, \dots, X_n]]$  like a sum of forms with respect to the usual degree. If  $f_r$  is a form of degree  $r$ , then  $\varphi(f_r) = t^r P$ , with  $P$  a polynomial in  $u$  and a finite number of elements  $u^{1/p^i}$ .

Let us take  $f, g \in k[[X_1, \dots, X_n]]$  such that  $g \neq 0$  and  $v(f/g) = 0$ . Then  $\varphi(f/g) = h_0 + th_1$ , where  $h_0$  is a rational fraction in  $u$  and a finite number of elements  $u^{1/p^i}$ . So  $h_0$  is algebraic over  $k(u)$ . Let us consider

$$P(u, Z) = c_0(u)Z^m + c_1(u)Z^{m-1} + \dots + b_{m-1}(u)Z + b_m(u)$$

a polynomial satisfied by  $h_0$ , where  $c_i \in k[u]$  for all  $i$  and  $c_0 \neq 0$ . Let  $\beta$  be the element

$$\beta = P\left(\frac{X_2}{X_1}, \frac{f}{g}\right) = c_0\left(\frac{X_2}{X_1}\right)\left(\frac{f}{g}\right)^m + \dots + c_n\left(\frac{X_2}{X_1}\right).$$

Then we have

$$\varphi(\beta) = c_0(u)(h_0 + th_1)^m + \dots + c_n(u),$$

so  $v(\beta) = \nu \circ \varphi(\beta) > 0$  and  $\beta \in \mathfrak{m}_v$ . Subsequently,

$$0 + \mathfrak{m}_v = \beta + \mathfrak{m}_v = P\left(\alpha, \frac{f}{g} + \mathfrak{m}_v\right).$$

This proves that  $f/g + \mathfrak{m}_v$  is an algebraic element over  $k(\alpha)$  and, a fortiori, the lemma.  $\square$

**Lemma 3.** *The dimension of a rank-one discrete valuation of the field  $k((X_1, \dots, X_n))$  is between 1 and  $n - 1$ .*

*Proof.* We know, after [2], that the dimension of a rank-one discrete valuation of  $k((X_1, \dots, X_n))$  is minor or equal than  $n - 1$ . So we have to prove that there exists a transcendental residue in  $\Delta_v$ .

Let us suppose that  $v(X_i) = n_i$  for all  $i = 1, \dots, n$ . Then the value of  $X_2^{n_1}/X_1^{n_2}$  is zero, so  $0 \neq (X_2^{n_1}/X_1^{n_2}) + \mathfrak{m}_v \in \Delta_v$ . If this residue lies in  $k$  then there exists  $a_{21} \in k$  such that

$$\frac{X_2^{n_1}}{X_1^{n_2}} + \mathfrak{m}_v = a_{21} + \mathfrak{m}_v.$$

This implies

$$\frac{X_2^{n_1}}{X_1^{n_2}} - a_{21} = \frac{X_2^{n_1} - a_{21}X_1^{n_2}}{X_1^{n_2}} \in \mathfrak{m}_v,$$

and then

$$v\left(\frac{X_2^{n_1} - a_{21}X_1^{n_2}}{X_1^{n_2}}\right) > 0.$$

So we have  $v(X_2^{n_1} - a_{21}X_1^{n_2}) = m_1 > n_1 n_2$ . Then

$$v\left(\frac{(X_2^{n_1} - a_{21}X_1^{n_2})^{n_1}}{X_1^{m_1}}\right) = 0.$$

If the residue of this element lies too in  $k$ , then there must exist  $a_{22} \in k$  such that  $v((X_2^{n_1} - a_{21}X_1^{n_2})^{n_1} - a_{22}X_1^{m_1}) = m_2 > n_1 m_1$ . We can repeat this operation.

The previous procedure is finite. If it didn't stop we would construct the power series

$$X_2 - \sum_{i=1}^{\infty} b_{2i} X_1^i$$

such that the sequence of partial sums has increasing values. Since  $\widehat{K}_n$  is a complete field, then this series amounts to zero in contradiction with  $X_1$  and  $X_2$  being formally independent. So the procedure must stop and there exists a transcendental element over  $k$  in  $\Delta_v$ .  $\square$

**Theorem 4.** *Let  $m$  be a fixed number between 1 and  $n - 1$ , then there exists a rank-one discrete valuation of  $k((X_1, \dots, X_n))$  of dimension  $m$ .*

*Proof.* Let us consider the one to one<sup>1</sup> homomorphism

$$\begin{aligned} \varphi: k[[X_1, \dots, X_n]] &\longrightarrow \overline{k(u)}[[t_1, \dots, t_m]] \\ X_1 &\longmapsto t_1 \\ X_2 &\longmapsto ut_1 \\ X_i &\longmapsto \begin{cases} t_i & \text{if } i \leq m + 1 \\ \sum_{j \geq 1} u^{1/p^j} t_1^j & \text{if } i > m + 1. \end{cases} \end{aligned}$$

We can take the valuation  $v := \nu \circ \varphi$ , with  $\nu$  the usual order function in  $\overline{k(u)}[[t_1, \dots, t_m]]$ . We know (lemma 2) that the residue  $X_2/X_1 + \mathfrak{m}_v$  is transcendental over  $k$ . Trivially the residue  $X_i/X_1 + \mathfrak{m}_v$  for all  $i = 3, \dots, m + 1$  are transcendental over  $k(X_2/X_1 + \mathfrak{m}_v, \dots, X_{i-1} + \mathfrak{m}_v)$  because  $t_i$  are formally independent variables. Any element  $f/g + \mathfrak{m}_v \in \Delta_v$  is algebraic over  $k(X_2/X_1 + \mathfrak{m}_v, \dots, X_{m+1} + \mathfrak{m}_v)$  by lemma 2. So the dimension of  $v$  is  $m$ .  $\square$

---

<sup>1</sup>The proof of injectivity parallels that of lemma 1.

## REFERENCES

1. E. Briaies, *Construcción explícita de valoraciones discretas de rango 1.*, Ph.D. thesis, Universidad de Sevilla, 1986.
2. ———, *Constructive theory of valuations.*, Comm. Algebra **17** (1989), no. 5, 1161–1177.
3. J. P. Serre, *Corps locaux.*, Hermann, Paris, 1968.

DEPARTAMENTO DE ÁLGEBRA, UNIVERSIDAD DE SEVILLA, ESPAÑA  
*E-mail address:* olalla@algebra.us.es