


PREPUBLICACIONES DEL DEPARTAMENTO DE ÁLGEBRA
DE LA UNIVERSIDAD DE SEVILLA

Monomial discrete valuations in $k[[\mathbf{X}]]$

Francisco Javier Herrera Govantes, Miguel Ángel Olalla Acosta, José Luis Vicente Córdoba

Prepublicación nº 23 (2 de octubre de 2003)



Departamento de Álgebra. Universidad de Sevilla

Monomial discrete valuations in $k[[\mathbf{X}]]$

Francisco Javier Herrera Govantes Miguel Ángel Olalla Acosta¹
José Luis Vicente Córdoba¹

*Departamento de Álgebra
Universidad de Sevilla
Apdo. 1160 – 41080 – Sevilla
SPAIN*

Abstract

We know that all the rank one discrete valuations of $k((X_1, X_2))$, centered in the ring $k[[X_1, X_2]]$, come from the usual order function [2,3], i.e. there exists a finite number of transformations such that we obtain a new field $k((Y_1, Y_2))$ where the lifting of v is a monomial valuation given by $v(Y_1) = v(Y_2) = 1$.

In this work we generalize this result to the rank m discrete valuation of $K = k((\mathbf{X}))$, centered in $R = k[[\mathbf{X}]]$. We prove that, if the dimension of v is $n - m$, the maximum since [1], then there exists an immediate extension L of K where the valuation is monomial. Therefore we give an explicit construction of the residue field of the valuation.

Key words: Valuation theory, formal power rings.

1 Preliminaries

Remark 1 *In this remark we remember some definitions and results given by I. Kaplansky in [4].*

Let K be a valued field, let v be the valuation. Let L be an extension of K and v' a valuation that extends v . We say that the extension $K \subseteq L$ is immediate if the values group and the residue field of v and v' are the same. We say that a valued field K is maximal if it doesn't admit proper immediate extensions.

Email addresses: `jherrera@algebra.us.es` (Francisco Javier Herrera Govantes), `miguelolalla@algebra.us.es` (Miguel Ángel Olalla Acosta), `jlvc@algebra.us.es` (José Luis Vicente Córdoba).

¹ Partially supported by BFM 2001-3207 (MCyT)

A well ordered set $\{a_i\} \subset K$, without last element in K , is called pseudo-convergent if

$$v(a_j - a_i) < v(a_k - a_j)$$

for all $i < j < k$. Easily we have that, if $\{a_i\}$ is pseudo-convergent, then $v(a_j - a_i) = v(a_{i+1} - a_i)$, $\forall i < j$. So we can use the abbreviation ω_i for $v(a_j - a_i)$, with $j > i$. Let us note that $\{\omega_i\}$ is an increasing set of elements of Γ .

An element $a \in K$ is called limit of the pseudo-convergent set $\{a_i\}$ if $v(a - a_i) = \omega_i$ for all i .

[4] prove the following results:

- If $K \subset L$ is an immediate extension, then every element of $L \setminus K$ is a limit of any pseudo-convergent subset of K without limit in K .
- A valued field K is maximal if and only if there exists a limit for all its pseudo-convergent subsets.

In [5], Krull shown the existence of, at least, one maximal immediate extension for all valued field. Therefore, if \widehat{K} is the completion of K by the valuation v and \widehat{v} is the unique valuation of \widehat{K} that extends v , then $K \subset \widehat{K}$ is an immediate extension and \widehat{K} is a maximal valued field.

Kaplansky shown the unicity of the maximal immediate extension of a valued field K that satisfy a certain condition, that he calls hypothesis A. In the case of zero characteristic this hypothesis A is empty.

Finally, [4] gives an useful structure theorem for all maximal valued field. If Δ is a field and Γ an ordered abelian group, the set of all formal series

$$\sum a_i t^{\alpha_i} \text{ con } a_i \in \Delta, \alpha_i \in \Gamma \text{ y } \{\alpha_i\} \text{ well ordered,}$$

is a field with the usual sum and times. We shall denote this field by $\Delta(t^\Gamma)$. In $\Delta(t^\Gamma)$ we can consider the valuation ν_t given by

$$\nu_t \left(\sum_{i \geq 1} a_i t^{\alpha_i} \right) = \alpha_1 \text{ with } a_1 \neq 0.$$

Krull shown in [5] that, with this valuation, $\Delta(t^\Gamma)$ is a maximal field.

Theorem 2 (Kaplansky, 1942) *Let K be a maximal valued field with values group Γ and residue field Δ that satisfies the hypothesis A. Then K is analytically isomorphic to the power series field $\Delta(t^\Gamma)$.*

1.1 Notation and definitions.

Let $K = k((\mathbf{X})) = k((X_1, \dots, X_n))$ be the quotient field of the formal power series ring $R = k[[\mathbf{X}]] = k[[X_1, \dots, X_n]]$.

Remark 3 1) We shall write all series $f \in R$ as $f = \sum_{A \in \mathbb{Z}_0^n} f_A \mathbf{X}^A$, where, if $A = (a_1, \dots, a_n)$, then \mathbf{X}^A means $X_1^{a_1} \dots X_n^{a_n}$. We shall say

$$\mathcal{E}(f) = \{A \in \mathbb{Z}_0^n \mid f_A \neq 0\}.$$

- 2) In \mathbb{Z}^m we'll consider the lexicographic order, it'll be denoted by \leq_{lex} . This is a total order for the group structure.
 3) Let $0 < m \leq n$ be an integer and

$$L = \{B_1, \dots, B_n\} \subset \mathbb{Z}_0^m \setminus \{0\}$$

such that L is a generator system of \mathbb{Z}^m . Each monomial \mathbf{X}^A of R has an element of \mathbb{Z}_0^m associated, that is called its L -degree, that is

$$\text{degree}_L(\mathbf{X}^A) = \sum_{i=1}^n a_i B_i, \quad A = (a_1, \dots, a_n).$$

Definition 4 Let $0 < m \leq n$ be an integer and

$$L = \{B_1, \dots, B_n\} \subset \mathbb{Z}_0^m \setminus \{0\}$$

such that L is a generator system of \mathbb{Z}^m . Let $v : R \rightarrow \mathbb{Z}^m \cup \{\infty\}$ be the function such that $v(0) = \infty$ and

$$v(f) = \min_{\leq_{\text{lex}}} \left\{ \text{degree}_L(\mathbf{X}^A) \mid A \in \mathcal{E}(f) \right\},$$

with $f \neq 0$. The extension of v to $K|k$, which values group is \mathbb{Z}^m , is a rank m discrete valuation, called monomial valuation associated to L .

Throughout this work, let v be a rank m discrete valuation of $K|k$ centered in R . Let R_v , \mathfrak{m}_v and $\Gamma = \mathbb{Z}^m$ be the ring, the maximal ideal and the values group of the valuation v , respectively. We'll denote by $\Delta_v = R_v/\mathfrak{m}_v$ to the residue field of v . Since [1] we know that the dimension of v (the transcendence degree of $k \subset \Delta_v$) is lesser or equal than $n - m$. We shall suppose that the dimension of v is the maximum, $n - m$.

So in this work valuation means rank m discrete valuation of $K|k$ centered in R and dimension $n - m$.

Remark 5 Let \widehat{K} be a maximal immediate extension of K . Since [4] we can suppose that \widehat{K} is the completion of K with respect of v . Let \widehat{v} the only extension of v to \widehat{K} . We know that there exists an analytic isomorphism of \widehat{K} in

$\Delta_v(t^\Gamma)$, so its restriction to the ring R gives an injective homomorphism

$$\begin{aligned}\varphi : R = k[[\mathbf{X}]] &\rightarrow \Delta_v(t^\Gamma) \\ X_i &\mapsto \sum_{j \geq 1} a_{i,j} t^{\alpha_{i,j}}\end{aligned}$$

with $a_{i,j} \in \Delta_v$ y $\{\alpha_{i,j}\} \subset \Gamma$ well ordered.

If we consider the extension of φ to the quotient field K and the valuation ν_t of $\Delta_v(t^\Gamma)$ previously defined, then $v = \nu_t \circ \varphi$.

The purpose of this work is to construct φ explicitly, in order to obtain a parametric equation of v and, in consequence, a construction of the residue field of v , as an extension of the field k .

Therefore we'll prove that for all valuation v of $K|k$, there exists an immediate extension $K \subset L = k((\mathbf{Y}))$ such that the valuation that extends v is monomial.

In other words: Any valuation v comes from a monomial valuation. This result generalizes the obtained in [2,3] for rank one discrete valuations of $k((X_1, X_2))$.

1.2 Monoidal Transformation and immediate extension.

Let v be a valuation of $K|k$. Let us consider the next monoidal transformation in K :

$$\begin{aligned}k((\mathbf{X})) &\rightarrow L = k((\mathbf{Y})) \\ X_i &\mapsto Y_i \text{ if } i \neq 2 \\ X_2 &\mapsto Y_2 Y_1\end{aligned}$$

with $v(X_2) >_{\text{lex}} v(X_1)$. Then we have the following theorem.

Theorem 6 *With these conditions, the extension $K \subset L$ is immediate.*

PROOF. Let us consider the rings $R = k[[\mathbf{X}]]$ and $S = k[[\mathbf{Y}]]$, and the diagram

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & \Delta_v(t^\Gamma) \\ \downarrow & \nearrow \psi & \\ S & & \end{array}$$

Where ψ is the natural extension of φ to L , i.e. $\psi(Y_2) = \varphi(X_2)/\varphi(X_1)$.

If ψ is injective, then $v' = \nu_t \circ \psi$ is a valuation of L that extends v and both has the same values group Γ and the same residue field Δ_v .

So we can suppose, by contradiction, that ψ is not injective. Let \mathfrak{p} be the implicit ideal $\ker(\psi)$ and L' the quotient field of the ring

$$\frac{k[[\mathbf{Y}]]}{\mathfrak{p}}.$$

Clearly the restriction of ψ to L' is injective and its composition with ν_t define a valuation, let us put w_1 . So the extension $K \subset L'$ is immediate.

Let w_2 be the valuation \mathfrak{p} -adic of L . This is a discrete rank one valuation.

The composition of both valuations, let us put w , is a discrete rank $m + 1$ valuation of $L|k$ whose residue field is Δ_v . Then we have

$$\text{rank}(w) + \dim(w) = n + 1 > \dim k[[\mathbf{Y}]].$$

Since Abhyankar's theorem ([1], Theorem 1, p. 330), we know

$$\text{rank}(w) + \dim(w) \leq \dim k[[\mathbf{Y}]],$$

so there is a contradiction.

Remark 7 *Here we are used the existence of such injective homomorphism, proved by Kaplansky [4]. Later we'll give an explicit construction of ψ . There isn't circular reasoning.*

The following example shows that the condition of maximal dimension of v is necessary.

Example 8 *Let $R = \mathbb{C}[[X_1, X_2, X_3]]$ and K its quotient field. Let us consider the injective homomorphism*

$$\begin{aligned} \varphi: R &\rightarrow \mathbb{C}(u)[[t]] \\ X_1 &\mapsto t \\ X_2 &\mapsto ut^2 \\ X_3 &\mapsto e^{ut} - 1. \end{aligned}$$

The composition of φ (really its extension to the quotient fields) with the usual order function ν_t of $\mathbb{C}(u)((t))$ is a rank one discrete valuation of K , named v . The dimension of v is 1, the transcendence degree of the extension $\mathbb{C} \subset \mathbb{C}(u)$.

If we make the monoidal transformation

$$\begin{aligned} R = \mathbb{C}[X_1, X_2, X_3] &\rightarrow S = \mathbb{C}[Y_1, Y_2, Y_3] \\ X_i &\mapsto Y_i \text{ if } i = 1, 3 \\ X_2 &\mapsto Y_2 Y_1 \end{aligned}$$

Then we have an homomorphism

$$\begin{aligned} \varphi: S &\rightarrow \mathbb{C}(u)[[t]] \\ X_1 &\mapsto t \\ X_2 &\mapsto ut \\ X_3 &\mapsto e^{ut} - 1. \end{aligned}$$

that is not injective.

2 Constructing Δ_v .

2.1 Basis of a subgroup of \mathbb{Z}^m .

There are well known procedures to compute a basis of a subgroup $\Gamma_0 \subset \mathbb{Z}^m$ knowing any set of generators. In this subsection we describe an algorithm that will be very useful in order to prepare the valuation.

Let $\{A_1, \dots, A_n\}$ be the set of generators of $\Gamma_0 \subset \mathbb{Z}^m$, with $A_i >_{\text{lex}} \mathbf{0}$ for all i . We can suppose, without loss of generality, that $A_i \leq_{\text{lex}} A_j \forall i < j$. Let $A = (a_{i,j}) \in M_{n \times m}$ be the matrix whose rows are the elements A_i .


We shall consider two transformations with the rows of the matrix A :

- (1) $F_{i,j}(q)$: To change the row i by itself plus q times the row j , with $q \in \mathbb{Z}$.
- (2) To interchange rows.

Clearly the group generated by the rows of the matrix A is equal to the group generated by the rows of any transformation of A .

Remark 9 *Algorithm.* The matrix A has the following echelon form

Let j be the first column different of $\mathbf{0}$ in A , let i be the first such that $a_{i,j} \neq 0$. Then we shall say that $a_{i,j}$ is a pivot.



How $A_i \leq_{\text{lex}} A_l \forall i < l$, clearly $a_{i,j} \leq a_{l,j} \forall i < l$. Let us put $a_{l,j} = q_l a_{i,j} + r_l$, by making the integer Euclidean division of $a_{l,j}$ by $a_{i,j}$.

We'll apply the following procedure with the first step, that is perfectly exportable to the other steps:

- I) For each row l , with $i < l \leq n$, we make the following transformation:
 - a) If $r_l \neq 0$ we do $F_{l,i}(-q_l)$. The new row l , that we denote by A_l again, is such that $A_l <_{\text{lex}} A_i$ and $0 < a_{l,j} < a_{i,j}$.
 - b) If $r_l = 0$ there are two possible situations:
 - 1) $A_l - q_l A_i >_{\text{lex}} \mathbf{0}$: In this case we make too the transformation $F_{l,i}(-q_l)$. The new row l is such that $A_l <_{\text{lex}} A_i$ and $a_{l,j} = 0$. So, after a reordering A_l raises a step.
 - 2) $A_l - q_l A_i \leq_{\text{lex}} \mathbf{0}$: Then we make the change $F_{l,i}(1 - q_l)$. The new row l , that we denote A_l again, is such that $A_l \leq_{\text{lex}} A_i$ and $m_{l,j} = m_{i,j}$. Let us remark that if $q_l = 1$ then $A_l = A_i$, because we have suppose at begin that $A_i \leq_{\text{lex}} A_l$.
- II) After these transformations we reorder the rows. If every row of the first step of the matrix are equal, then we raise a step and we begin with the procedure. In other case, we apply again this algorithm.

Clearly we are doing the Euclidean algorithm in order to compute the maximal common divisor of the elements $\{a_{i,j}, \dots, a_{n,j}\}$. Let p_j be the maximal common divisor. By a finite number of transformations we obtain a transformed matrix, that we denote again by A , such that the pivot is equal to p_j and the last one divides all $a_{l,j}$ with $l \geq i$.

We have just arrived to situation I.b), so, by a finite number of transformations, necessarily we must obtain a new matrix A where all the rows down the pivot $a_{i,j} = p_j$ are equals to A_i .

Hence, by applying this algorithm for each step of the matrix, we obtain a matrix B with only s different rows B_{i_1}, \dots, B_{i_s} whose pivots are p_{j_1}, \dots, p_{j_s} . Clearly Γ_0 is isomorphic to $p_{j_1}\mathbb{Z} \times \dots \times p_{j_s}\mathbb{Z}$ and $\{B_{i_1}, \dots, B_{i_s}\}$ is a basis of Γ_0 .

The algorithm described in this section allows us to prove the following lemma, that we are going to use frequently for preparing our valuation conveniently.

Lemma 10 *With the usual conditions over $K = k(\mathbf{X})$ and v , rank m dis-*

crete valuation, if Γ_0 is the subgroup generated by the values of the elements X_i , then we can find, by a finite number of monoidal transformations and interchanges of variables, an immediate extension $L = k(\mathbf{Y})$ of K such that each Y_i has the value in a basis $\{B_1, \dots, B_s\}$ of Γ_0 .

PROOF. We have to apply the precedent algorithm to the matrix of the values $v(X_i)$. Where $F_{l,i}(-q_l)$ means “to apply q_l monoidal transformations such that $X_l \mapsto Y_l Y_i$ ”, and reorder rows means “to reorder variables according to its values”.

2.2 Preparing v .

As usually, we begin with a rank $m \leq n$ discrete valuation v of $K|k$, centered in the ring $R = k[\mathbf{X}]$. Let $\Gamma = \mathbb{Z}^m$, R_v and \mathfrak{m}_v be the values group, the ring and the maximal ideal of the valuation. v , respectively. We shall denote, as usual, by Δ_v to the residue field of the valuation.

We'll suppose that the extension $k \subset \Delta_v$ is transcendental pure of degree $\dim v = n - m$.

First, we apply the lemma 10 to obtain an immediate extension $L = k(\mathbf{Y})$ of K such that the set of values of the elements Y_i is a basis $\{B_1, \dots, B_s\}$ of the subgroup Γ_0 generated by the values of the elements X_i .

By convenience we reorder the elements Y_i in such way that the first s elements takes all the values of the basis (i.e. $v(Y_i) = B_i$ for $i = 1, \dots, s$).

Remark 11 Let $A \in \Gamma_0$ be such that $A = r_1 B_1 + \dots + r_s B_s$, then we shall denote by $R_A = (r_1, \dots, r_s, 0, \dots, 0)$ to the n -uple such that $v(\mathbf{Y}^{R_A}) = A$.

2.3 The first transcendental residue.

Our purpose is to give an explicit description of the injective homomorphism $\psi: k[\mathbf{Y}] \rightarrow \Delta_v(t^\Gamma)$, such that $v = \nu_t \circ \psi$. In order not to complicate the exposition of this construction, we are supposing that all the residues are in k or they are transcendental over the ground field. This condition seems too strong, but after the proof of theorem 16 we'll explain why this situation is really close to the general one.

1) For the first elements we put

$$\psi(Y_i) = t^{B_i} \quad i = 1, \dots, s.$$

2) Let us take Y_{s+1} the first element whose value is a linear combination of the elements B_i (in fact it must be equal to some B_i). Let us suppose that $v(Y_{s+1}) = B_{s+1,1}$, then we have two possibilities:

- a) For all $\alpha \in k$, $v(Y_{s+1} + \alpha \mathbf{Y}^{R_{B_{s+1,1}}}}) = B_{s+1,1}$. This fact means that the residue of $Y_{s+1}/\mathbf{Y}^{R_{B_{s+1,1}}}$ in Δ_v is transcendental over k . Let us put

$$u_{s+1} = \frac{Y_{s+1}}{\mathbf{Y}^{R_{B_{s+1,1}}}} + \mathfrak{m}_v$$

and $\psi(Y_{s+1}) = u_{s+1}t^{B_{s+1,1}}$.

- b) There exists $\alpha \in k$ such that $v(Y_{s+1} + \alpha \mathbf{Y}^{R_{B_{s+1,1}}}) >_{\text{lex}} B_{s+1,1}$. So the residue of $Y_{s+1}/\mathbf{Y}^{R_{B_{s+1,1}}}$ in Δ_v is in k . Let us put $\alpha_{s+1,1} = \alpha$ and $v(Y_{s+1} + \alpha_{s+1,1} \mathbf{Y}^{R_{B_{s+1,1}}}) = B_{s+1,2} > B_{s+1,1}$. There are two possibilities again:

- i) The new value $B_{s+1,2} \notin \Gamma_0$. In this case, we make the change $Z_{s+1} = Y_{s+1} + \mathbf{Y}^{R_{B_{s+1,1}}}$, $Z_i = Y_i \forall i \neq s+1$ and go back to the beginning of the procedure by preparing the new valuation of $k((\mathbf{Z}))$ with the lemma 10.
- ii) The value $B_{s+1,2} \in \Gamma_0$. If there exists $\alpha_{s+1,2}$ such that

$$v\left(Y_{s+1} + \alpha_{s+1,1} \mathbf{Y}^{R_{B_{s+1,1}}} + \alpha_{s+1,2} \mathbf{Y}^{R_{B_{s+1,2}}}\right) = B_{s+1,3} >_{\text{lex}} B_{s+1,2},$$

then we ask again if $B_{s+1,3} \in \Gamma_0$. In the affirmative case, if there exists $\alpha_{s+1,3} \in k$ such that

$$v\left(Y_{s+1} + \alpha_{s+1,1} \mathbf{Y}^{R_{B_{s+1,1}}} + \alpha_{s+1,2} \mathbf{Y}^{R_{B_{s+1,2}}} + \alpha_{s+1,3} \mathbf{Y}^{R_{B_{s+1,3}}}\right) =$$

$$B_{s+1,4} >_{\text{lex}} B_{s+1,3},$$

we go back to the beginning of the procedure.

We continue this procedure until we find a value $B_{s+1,l} \notin \Gamma_0$. If we cannot, there does not exist $\alpha_{s+1,l} \in k$ such that

$$v\left(Y_{s+1} + \sum_{k=1}^l \alpha_{s+1,k} \mathbf{Y}^{R_{B_{s+1,k}}}\right) = B_{s+1,l+1} > B_{s+1,l}.$$

In the first case we make the change

$$Z_{s+1} = Y_{s+1} + \sum_{k=1}^{l-1} \alpha_{s+1,k} \mathbf{Y}^{R_{B_{s+1,k}}}, \quad Z_i = X_i \quad \forall i \neq s+1$$

and move to the lemma 10. In the second case we have that the residue

$$u_{s+1} = \frac{Y_{s+1} + \sum_{k=1}^{l-1} \alpha_{s+1,k} \mathbf{Y}^{R_{B_{s+1,k}}}}{\mathbf{Y}^{R_{B_{s+1,l}}}} + \mathfrak{m}_v$$

is transcendental over k . In this case we put

$$\psi(Y_{s+1}) = \sum_{k=1}^{l-1} \alpha_{s+1,k} t^{B_{s+1,k}} + u_{s+1} t^{B_{s+1,l}}.$$

We have to prove that this procedure ends up finding a new value that it is not in Γ or a transcendental residue.

Lemma 12 *The situation described in 2.b.i) only occurs a finite number of times.*

PROOF. There are two possible situations to find an element of value $B \notin \Gamma_0$:

1) The value B is not a rational linear combination of the elements of Γ_0 . In this case, the rank of the new subgroup of \mathbb{Z}^m increases by 1. Trivially this fact only occurs a finite number of times.

2) The new value B is a non integer rational linear combination of the elements of the basis of Γ_0 . Let $\{p_1, \dots, p_s\}$ be the pivots that appears in the construction of the basis of Γ_0 , let $\{q_1, \dots, q_s\}$ be the ones of the new subgroup, Γ_1 . As $\Gamma_0 \subset \Gamma_1$, then $q_i \leq p_i \forall i$ and, at least, one inequality is strict. As the pivotes are greater or equal than 1, this only occurs a finite number of times.

Remark 13 *Let us suppose that we have a pseudo convergent set $\{f_j\}$ of elements of a valued field K , with value group \mathbb{Z}^m with lexicographic order. The set of values $\{\omega_j\}$ of the elements f_j is a strictly increasing sequence of elements of the group \mathbb{Z}^m . Let us suppose that the set $\{\omega_j\}$ is bounded. Then, from an index j sufficiently big, we have*

$$\omega_j = (a_1, \dots, a_l, a_{l+1,j}, \dots, a_{m,j}),$$

in such way that the first l coordinates of the values ω_j are stabilized. Let us suppose that l is the greatest integer between 1 and m such that this fact occurs.

Let f be a limit of $\{f_j\}$ and $\omega = v(f)$ its value. Then, if $\omega = (b_1, \dots, b_n)$, as $\omega >_{\text{lex}} \omega_j$ for all j , then there exists an $l_0 \leq l$ such that $b_i = a_i$ for all $i = 1, \dots, l_0 - 1$ and $b_{l_0} > a_{l_0}$.

Lemma 14 *The procedure described in the situation 2.b.ii) finds a value that is not in Γ_0 or a transcendental residue over k .*

PROOF. If the procedure is not finite, then we have a set of elements $\{f_j\}_{j \geq 1}$ such that

$$f_j = Y_{s+1} + \sum_{l=1}^{j-1} \alpha_{s+1,l} \mathbf{Y}^{R_{B_{s+1},l}}.$$

So $v(f_{j+1}) = B_{s+1,j+1} >_{\text{lex}} v(f_j) = B_{s+1,j}$ for all j .

Therefore, if $i < j < k$, we have

$$v(f_j - f_i) = B_{s+1,i} <_{\text{lex}} v(f_k - f_j) = B_{s+1,j},$$

so $\{f_j\}$ is a pseudo-convergent set.

Let us suppose that the set of values $\{B_{s+1,j}\}$ is bounded. Like in the remark 13, let us suppose that, since an index j sufficiently great, the first l coordinates of the values are stabilized, where l is the greatest integer between 1 and m such that this fact occurs. Let us put

$$v(f_j) = (a_1, \dots, a_l, a_{l+1,j}, \dots, a_{m,j}).$$

Again by remark 13, we know that any limit of $\{f_j\}$ is such that its value is something like

$$(a_1, \dots, a_k, b_{k+1}, \dots, b_m),$$

with $k < l$ and $b_{k+1} > a_{k+1}$. Let us take the series

$$Y_{s+1} + \sum_{j=1}^{\infty} \alpha_{s+1,j} \mathbf{Y}^{R_{B_{s+1,j}}}.$$

As limit, by convenience we put

$$g_1 = \sum_{j=1}^{\infty} \alpha_{s+1,j} \mathbf{Y}^{R_{B_{s+1,j}}}$$

and $v(Y_{s+1} + g_1) = B_{s+1,1}^1$.

If $B_{s+1,1}^1 \notin \Gamma_0$, then we have finished. In other case, if there exists $\alpha_{s+1,1}^1$ such that

$$v\left(Y_{s+1} + g_1 + \alpha_{s+1,1}^1 \mathbf{Y}^{R_{B_{s+1,1}^1}}\right) = B_{s+1,2}^1 >_{\text{lex}} B_{s+1,1}^1,$$

then we continue with our procedure. If it is infinite again and the values contained are bounded, then we shall have a pseudo-convergent set with limit

$$Y_{s+1} + g_1 + \sum_{j=1}^{\infty} \alpha_{s+1,j}^1 \mathbf{Y}^{R_{B_{s+1,j}^1}}.$$

Again by convenience we put

$$g_2 = \sum_{j=1}^{\infty} \alpha_{s+1,j}^1 \mathbf{Y}^{R_{B_{s+1,j}^1}}$$

and $v(Y_{s+1} + g_1 + g_2) = B_{s+1,1}^2 >_{\text{lex}} B_{s+1,1}^1$.

If this procedure does not find an element not in Γ_0 or a transcendental residue, we have a series

$$Y_{s+1} + g_1 + g_2 + g_3 + \dots$$

such that the set of the partial sums

$$\left\{ Y_{s+1} + \sum_{i=1}^j g_i \right\}_{j \geq 1}$$

is pseudo-convergent.

If the values of this set are bounded again, then, from one j sufficiently great, the first k coordinates of each value are stabilized, where k is the greatest integer between 1 and m such that this fact occurs. As all the values of this set are greater than the first limit we have found, then it must be $k < l$.

Therefore, if we don't find a transcendental residue, after an infinite procedure of calculation of limits of pseudo-convergent sets, the values, if they are bounded, become stable in lesser and lesser coordinates.

So this procedure must end finding an element that is not in Γ_0 or a new transcendental residue. The opposite fact is equivalent to construct a series

$$f = Y_{s+1} + g(Y_1, \dots, Y_s) \in k((\mathbf{Y}))$$

such that the set of values of the partial sums is not a bounded strictly increasing sequence. This means that $v(f) = \infty$. So $\psi(f) = 0$, but this is a contradiction because ψ is injective.

So, at the end of this procedure, we shall have a series $f = Y_{s+1} + g(Y_1, \dots, Y_s) \in k((\mathbf{Y}))$ such that $v(f) = B_{s+1}$ and, either $B_{s+1} \notin \Gamma_0$, or there does not exist α such that $v(f + \alpha \mathbf{Y}^{R_{B_{s+1}}}) >_{\text{lex}} B_{s+1}$. In this last case we take

$$u_{s+1} = \frac{f}{\mathbf{Y}^{R_{B_{s+1}}}} + \mathbf{m}_v$$

and put

$$\psi(Y_{s+1}) = g(t^{B_1}, \dots, t^{B_s}) + u_{s+1} t^{B_{s+1}}.$$

2.4 Construction of Δ_v .

We can suppose that we have an immediate extension $L = k((\mathbf{Y}))$ of K such that:

1) $\{B_1, \dots, B_m\}$ is a basis of $\Gamma = \mathbb{Z}^m$. This means that the situation 2.b.i) is not going to appear any more and all the values that we find are integer linear combination of the basis.

2) We have applied the described procedure with the first $k > m$ elements Y_i obtaining:

$$\begin{aligned}\psi(Y_i) &= t^{B_i}, \quad i = 1, \dots, m \\ \psi(Y_j) &= \sum_{l \geq 1} \alpha_{j,l} t^{B_{j,l}} + u_j t^{B_j}, \quad j = m+1, \dots, k\end{aligned}$$

such that $B_j >_{\text{lex}} B_{j,l} >_{\text{lex}} B_{j,l-1}$ y $\{\alpha_{j,l}\} \subset \Delta_{j-1} = k(u_{m+1}, \dots, u_{j-1}) \subset \Delta_v$.

Our purpose is to describe the procedure with the variable Y_{k+1} , that is analogous to the described previously to find the first transcendental residue.

Let $v(Y_{k+1}) = B_{k+1,1}$, there exists $\alpha_{k+1,1} \in \Delta_k$ such that the value $v(Y_{k+1} + \alpha_{k+1,1} \mathbf{Y}^{R_{B_{k+1,1}}}) = B_{k+1,2} >_{\text{lex}} B_{k+1,1}$?

1) If the answer is affirmative, then we pose the question for the new element $Y_{k+1} + \alpha_{k+1,1} \mathbf{Y}^{R_{B_{k+1,1}}}$.

2) If it is negative then we put

$$\psi(Y_{k+1}) = u_{k+1} t^{B_{k+1,1}}$$

and go to the following variable.

Lemma 15 *This procedure ends finding a new transcendental residue, eventually after an infinite number of calculations.*

PROOF. The reasoning is the same that in lemma 14, if there is not such transcendental residue, then we can construct a pseudo-convergent set that has a limit

$$f = Y_{k+1} + g(Y_1, \dots, Y_m) \in k((\mathbf{Y})),$$

of value ∞ . This implies that $\psi(f) = 0$, and this is a contradiction

With this procedure we prove the following theorem:

Theorem 16 *The residue field of v is $k(u_{m+1}, \dots, u_n)$.*

PROOF. The constructions described in this work permit us to find one immediate extension $L = k((\mathbf{Y}))$ of K such that the valuation that extends v to the field L is defined by the composition of

$$\begin{aligned}\psi : k((\mathbf{Y})) &\rightarrow k(u_{m+1}, \dots, u_n)(t^\Gamma) \\ Y_i &\mapsto t^{B_i}, \quad i = 1, \dots, m \\ Y_j &\mapsto \sum_{k \geq 1} \alpha_{j,k} t^{B_{j,k}} + u_j t^{B_j}, \quad j = m+1, \dots, n\end{aligned}$$

with the valuation ν_t of $k(u_{m+1}, \dots, u_n)(t^\Gamma)$. Clearly the residue field of this valuation is $k(u_{m+1}, \dots, u_n)$.

Remark 17 *The general case: considering algebraic residues. If we find non-trivial algebraic residues, then the procedure does not change, and we obtain the residue field*

$$\Delta_v = k(\zeta_{m+1}, \dots, \zeta_n),$$

where ζ_i is a collection, eventually infinite, of algebraic residues $\{\alpha_{i,k}\}$ and one transcendental residue u_i .

All of these algebraic extensions are finite; in other case we can do the same as the proof of theorem 6 to obtain a valuation with dimension greater than $n - m$.

It could be another problem in the general case, because in fact we are constructing a representant field of Δ_v as an intermediate extension of $k \subset \sigma(\Delta_v) \subset K$, where σ is a section of the natural homomorphism $R_v \rightarrow \Delta_v$. As the rings of power series are completes, we can apply Hensel's lemma to know that we can find an algebraic representant $\alpha \in K$ for every algebraic residue $\alpha + \mathfrak{m}_v$.

3 Monomial valuations.

Although the number of operations to calculate ψ explicitly with the previous procedure is infinite, the number of monoidal transformations, coordinates changes and interchanges of variables that transform K in L is finite, because these only appear when we find a value $B \notin \Gamma_0$ and this fact occurs a finite number of times. Following the trace of these transformations we obtain a map $\varphi : K \rightarrow \Delta_v(t^\Gamma)$ that is the restriction of ψ and parametrizes v . In a similar way we can obtain some representatives of every residue u_i depending on \mathbf{X} .

Finally we give a theorem that generalizes the results obtained in [2,3] for rank one discrete valuations of $k((X_1, X_2))$ centered in $k[[X_1, X_2]]$.

Theorem 18 *For all rank m discrete valuation v of $K|k$, with dimension $n - m$ and centered in R , there exists an immediate extension $L = k((\mathbf{Z}))$ of K such that the lifting of v to L is monomial.*

PROOF. In fact, by means of the previous procedure we obtain an immediate

extension $k((\mathbf{Y}))$ of K such that the lifting of v is $v = \nu_t \circ \psi$, with

$$\begin{aligned}\psi : k((\mathbf{Y})) &\rightarrow k(u_{m+1}, \dots, u_n)(t^\Gamma) \\ Y_i &\mapsto t^{B_i}, \quad i = 1, \dots, m \\ Y_j &\mapsto \sum_{k \geq 1} \alpha_{j,k} t^{B_{j,k}} + u_j t^{B_j}, \quad j = m+1, \dots, n.\end{aligned}$$

This valuation is not monomial, but if we make the change

$$\begin{aligned}k((\mathbf{Y})) &\rightarrow k((\mathbf{Z})) \\ Y_i &\mapsto Z_i, \quad i = 1, \dots, m \\ Y_j &\mapsto Z_j - \sum_{k \geq 1} \alpha_{j,k} \mathbf{Z}^{R_{B_{j,k}}}, \quad j = m+1, \dots, n,\end{aligned}$$

then the natural extension of ψ to $k((\mathbf{Z}))$ is

$$\begin{aligned}\phi : k((\mathbf{Z})) &\rightarrow k(u_{m+1}, \dots, u_n)(t^\Gamma) \\ Y_i &\mapsto t^{B_i}, \quad i = 1, \dots, m \\ Y_j &\mapsto u_j t^{B_j}, \quad j = m+1, \dots, n\end{aligned}$$

and the valuation $v = \nu_t \circ \phi$ is monomial.

Remark 19 *In the general case, considering algebraic residues, we obtain an immediate extension $L = k'((\mathbf{Z}))$, where k' is an extension of k contained in $\sigma(\Delta_v)$. The lifting of v in L is a monomial valuation.*

Example 20 *Let $R = k[[X_1, X_2, X_3, X_4]]$ be the power series ring, with $k = \mathbb{Z}/\mathbb{Z}_5$. Let K be the quotient field. Let us consider the series $W_1 = X_4 - \sum_{i \geq 1} X_3^{3i}$ and $W_2 = X_2 - \sum_{i \geq 1} X_1^i$. Let us consider the following valuations:*

v_1 : (W_1) -adic valuation of K ,

v_2 : (W_2) -adic valuation of $K/(W_1)$ and

v_3 : the monomial valuation of $K/(W_1, W_2)$ defined by the values $v_3(X_1) = v_3(X_3) = 1$.

Finally, let v be the composite valuation $v = v_3 \circ v_2 \circ v_1$. It is a rank 3 discrete valuation of $K|k$, such that:

$$\begin{aligned}v(X_1) &= v(X_3) = (0, 0, 1) \\ v(W_2) &= (0, 1, 0) \\ v(W_1) &= (1, 0, 0),\end{aligned}$$

wherefrom

$$v(X_2) = v(W_2 + \sum_{i \geq 1} X_1^i) = (0, 0, 1)$$

$$v(X_4) = v(W_1 + \sum_{i \geq 1} X_3^{3i}) = (0, 0, 3).$$

So the values of the variables generate the subgroup $\Gamma_0 = \{0\} \times \{0\} \times \mathbb{Z}$. From lemma 10 we know that the transformation $X_4 \mapsto Y_4 Y_1^2$ takes K in to an immediate extension $L = k((Y_1, Y_2, Y_3, Y_4))$, such that $v(Y_i) = (0, 0, 1)$, $i = 1, 2, 3, 4$. We shall apply the given procedure with this example to obtain a parametrization, ψ , of v .

Let us put $\psi(Y_1) = t^{(0,0,1)}$. We shall begin with the variable Y_2 , as $v(Y_2) = (0, 0, 1)$, and we ask if there exists $\alpha \in k$ such that $v(Y_2 + \alpha Y_1) >_{\text{lex}} (0, 0, 1)$. By the construction of v , we know that $v(Y_2 - Y_1) = v(X_2 - X_1) = (0, 0, 2) >_{\text{lex}} (0, 0, 1)$. Following the procedure, we find that

$$v\left(Y_2 - \sum_{i=1}^l i Y_1^i\right) = (0, 0, l+1) >_{\text{lex}} (0, 0, l).$$

So we have a pseudo-convergent set $\{f_l\}$, with $f_l = Y_2 - \sum_{i=1}^l i Y_1^i$. A limit of this set is

$$f_\infty = Y_2 - \sum_{i \geq 1} i Y_1^i.$$

As $v(f_\infty) = v(W_2) = (0, 1, 0) \notin \Gamma_0$, we put

$$\psi(Y_2) = \sum_{i \geq 1} i t^{(0,0,i)} + t^{(0,1,0)}.$$

We continue with Y_3 , now $\Gamma_0 = \{0\} \times \mathbb{Z} \times \mathbb{Z}$. Since it does not exist $\alpha \in k$ such that $v(Y_3 + \alpha Y_1) >_{\text{lex}} (0, 0, 1)$, we put

$$\psi(Y_3) = u_3 t^{(0,0,1)}.$$

Let $\Delta_3 = k(u_3)$, with $u_3 = Y_3/Y_1 + \mathfrak{m}_v$. Let us go to the variable $Y_4 = X_4/X_1^2$, $v(Y_4) = (0, 0, 1)$. By the construction of v , we know that

$$v\left(Y_4 - u_3^3 Y_1\right) = v\left(\frac{X_4 - X_3^3}{X_1^2}\right) = (0, 0, 4) >_{\text{lex}} (0, 0, 1).$$

Following the procedure we have

$$v\left(Y_4 - \sum_{i=1}^l u_3^{3i} Y_1^{3i-2}\right) = v\left(\frac{X_4 - \sum_{i=1}^l X_3^{3i}}{X_1^2}\right) = (0, 0, 3i+1).$$

Wherefrom we come to have a pseudo-convergent set $\{g_l\}$, with $g_l = Y_4 -$

$\sum_{i=1}^l u_3^{3i} Y_1^{3i-2}$. A limit of this set is

$$g_\infty = Y_4 - \sum_{i \geq 1} u_3^{3i} Y_1^{3i-2}$$

and $v(g_\infty) = v(W_1/X_1^2) = (1, 0, -2) \notin \Gamma_0$. We put

$$\psi(Y_4) = \sum_{i \geq 1} u_3^{3i} t^{(0,0,3i-2)} + t^{(1,0,-2)}.$$

Therefore we have $\Delta_v = k(X_3/X_1 + \mathfrak{m}_v)$ and

$$\begin{cases} \psi(X_1) = t^{(0,0,1)} \\ \psi(X_2) = \sum_{i \geq 1} i t^{(0,0,i)} + t^{(0,1,0)} \\ \psi(X_3) = u_3 t^{(0,0,1)} \\ \psi(X_4) = \sum_{i \geq 1} u_3^{3i} t^{(0,0,3i)} + t^{(1,0,0)}. \end{cases}$$

This way, making the substitution $X_1 = Z_1$, $X_2 = Z_2 + \sum i Z_1^i$, $X_3 = Z_3$ y $X_4 = Z_4 + \sum Z_3^{3i}$, we have an immediate extension $M = k((Z_1, Z_2, Z_3, Z_4))$ of K . The valuation that extends v to M is the monomial valuation defined by $v(Z_1) = v(Z_3) = (0, 0, 1)$, $v(Z_2) = (0, 1, 0)$ and $v(Z_4) = (1, 0, 0)$

References

- [1] S.S. Abhyankar, On the valuations centered in a local domain, *Amer. J. Math.* **78** (1956), 321–348.
- [2] E. Briaies, Constructive theory of valuations, *Commun. Algebra* **17** (1989), no. 5, 1161–1177.
- [3] E. Briaies and F.J. Herrera, Construcción explícita de las valoraciones de un anillo de series formales en dos variables, *Actas X Jornadas Hispano-Lusas* (Murcia), vol. II, 1985, pp. 1–10.
- [4] I. Kaplansky, Maximal fields with valuations, *Duke Math. J.* **9** (1942), 303–321.
- [5] W. Krull, Allgemeine bewertungstheorie, *J. Reine Angew. Math.* **167** (1931), 160–196.